

Realised power variation and stochastic volatility models

OLE E. BARNDORFF-NIELSEN

*The Centre for Mathematical Physics and Stochastics (MaPhySto),
University of Aarhus, Ny Munkegade, DK-8000 Aarhus C, Denmark.*

oebn@mi.aau.dk

NEIL SHEPHARD

Nuffield College, Oxford OX1 1NF, UK

neil.shephard@nuf.ox.ac.uk

June 26, 2002

Abstract

Limit distribution results on realised power variation, that is sums of absolute powers of increments of a process, are derived for certain types of semimartingale with continuous local martingale component, in particular for a class of flexible stochastic volatility models. The theory covers, for example, the cases of realised volatility and realised absolute variation. Such results should be helpful in, for example, the analysis of volatility models using high frequency information.

Some key words: Absolute returns; Mixed asymptotic normality; Realised volatility; p -variation; Quadratic variation; Semimartingale.

1 Introduction

Stochastic volatility processes play an important role in financial economics, generalising Brownian motion to allow the scale of the increments (or returns in economics) to change through time in a stochastic manner. We show such intermittency can be coherently measured using sums of absolute powers of increments, which we name realised power variation. This paper derives limit theorems for these measures, over a fixed interval of time, as the number of high frequency increments goes off to infinity.

A referee has drawn our attention to an unpublished thesis by Etienne Becker (1998) that develops a number of results that are closely related to those of the present paper. We outline the relation to Becker's work in the concluding Section of this paper.

This paper has six other sections. In Section 2 we establish our idea of realised power variation as well as define the regularity assumptions we need to derive our limit theorems. Section 3 contains our main results, while the proofs of them are given in Section 4. Section 5 gives some examples of the processes covered by our theory, while a Monte Carlo experiment to assess the accuracy of our limit theory approximations is conducted in Section 6. Finally,

Section 7 gives some concluding remarks including a discussion of the use of these ideas in other areas of study, for instance turbulence and image analysis.

2 Models, notation and regularity conditions

We first introduce some notation for realised power variation quantities of an arbitrary semimartingale x . Let δ be positive real and, for any $t \geq 0$, define

$$x_\delta(t) = x(\lfloor t/\delta \rfloor \delta),$$

where $\lfloor a \rfloor$ for any real number a denotes the largest integer less than or equal to a . The process $x_\delta(t)$ is a discrete approximation to $x(t)$. Further, for r positive real we define the *realised power variation of order r* ¹ or *realised r -tic variation* of $x_\delta(t)$ as

$$\begin{aligned} [x_\delta]^{[r]}(t) &= \sum_{j=1}^M |x_\delta(j\delta) - x_\delta((j-1)\delta)|^r \\ &= \sum_{j=1}^M |x(j\delta) - x((j-1)\delta)|^r \end{aligned} \quad (1)$$

where $M = M(t) = \lfloor t/\delta \rfloor$. Then, in particular, for $M \rightarrow \infty$, *realised quadratic variation*

$$[x_\delta]^{[2]}(t) \xrightarrow{p} [x](t),$$

where $[x]$ is the *quadratic variation* process of the semimartingale x . Note also that,

$$[x_\delta]^{[2]} = [x_\delta].$$

Henceforth, for simplicity of exposition, we consider only a fixed t and take δ so that $M = \lfloor t/\delta \rfloor$ is an integer (and then $\delta M = t$). However, our results can undoubtedly be generalised to provide functional limit statements, and we hope to discuss that elsewhere.

Our detailed results will be established for the stochastic volatility (SV) model where basic Brownian motion is generalised to allow the volatility term to vary over time and there to be a rather general drift. Then the y^* follows

$$y^*(t) = \alpha^*(t) + \int_0^t \sigma(s)dw(s), \quad t \geq 0, \quad (2)$$

¹The similarly named p -variation, $0 < p < \infty$, of a real-valued function f on $[a, b]$ is defined as

$$\sup_{\kappa} \sum |f(x_i) - f(x_{i-1})|^p,$$

where the supremum is taken over all subdivisions κ of $[a, b]$. If this function is finite then f is said to have bounded p -variation on $[a, b]$. The case of $p = 1$ gives the usual definition of bounded variation.

This concept has been studied recently in the probability literature. See the work of, for example, Lyons (1994) and Mikosch and Norvaiša (2000).

where $\sigma > 0$ and α^* are assumed to be stochastically independent of the standard Brownian motion w . Throughout this paper we will assume that the processes $\tau = \sigma^2$ and α^* predictable and pathwise locally Riemann integrable (hence, in particular, pathwise locally bounded). Thus y^* is a semimartingale, more precisely a Brownian semimartingale. We call σ the *spot volatility* process and α^* the *mean* or *risk premium* process. (For some general information on processes y^* of this type from the viewpoint of financial econometrics, see for example Ghysels, Harvey, and Renault (1996) and Barndorff-Nielsen and Shephard (2001)). By allowing the spot volatility to be random and serially dependent, this model will imply its increments will exhibit volatility clustering and have unconditional distributions which are fat tailed. This allows it to be used in finance and econometrics as a model for log-prices. In turn, this provides the basis for option pricing models which overcome some of the major failings in the Black-Scholes option pricing approach. Leading references in this regard include Hull and White (1987), Heston (1993) and Renault (1997). See also the recent work of Nicolato and Venardos (2001).

For the price process (2) the *realised power variation of order r* of y^* is, at time t and discretisation δ , $[y_\delta^*]^{[r]}(t)$. Letting

$$y_j(t) = y^*(j\delta) - y^*((j-1)\delta)$$

we have that

$$[y_\delta^*]^{[r]}(t) = \sum_{j=1}^M |y_j(t)|^r.$$

We use the notation $\tau(t) = \sigma^2(t)$ and

$$\tau^*(t) = \int_0^t \tau(s) ds$$

and, more generally, we consider the *integrated power volatility of order r*

$$\tau^{r*}(t) = \int_0^t \tau^r(s) ds.$$

Throughout the following, r denotes a positive number. Moreover we shall refer to the following conditions on the volatility and mean processes:

(V) The volatility process $\tau = \sigma^2$ is (pathwise) locally bounded away from 0 and has, moreover, the property

$$\lim_{\delta \downarrow 0} \delta^{1/2} \sum_{j=1}^M |\tau^r(\eta_j) - \tau^r(\xi_j)| = 0 \tag{3}$$

for some $r > 0$ (equivalently for all $r > 0$)² and for any $\xi_j = \xi_j(\delta)$ and $\eta_j = \eta_j(\delta)$ such that

$$0 \leq \xi_1 \leq \eta_1 \leq \delta \leq \xi_2 \leq \eta_2 \leq 2\delta \leq \dots \leq \xi_j \leq \eta_j \leq M\delta = t.$$

(M) The mean process α^* satisfies (pathwise)³

$$\overline{\lim}_{\delta \downarrow 0} \max_{1 \leq j \leq M} \delta^{-1} |\alpha^*(j\delta) - \alpha^*((j-1)\delta)| < \infty. \quad (4)$$

These regularity conditions are quite mild.⁴ Of some special interest are cases where α^* is of the form

$$\alpha^*(t) = \int_0^t g(\sigma(s)) ds,$$

for g a smooth function. Then regularity of τ will imply regularity of α^* .

Note that the assumptions allow the spot volatility to have, for example, deterministic diurnal effects, jumps, long memory, no unconditional mean or to be non-stationary.

On the other hand, models for τ like the Cox-Ingersoll-Ross process, or more generally the constant elasticity of variance process, do not satisfy (V). The same limit results do, however, hold also for such processes (at least for $r < 3$) as verified in a forthcoming paper by Ole Barndorff-Nielsen, Svend Erik Graversen and Neil Shephard.

3 Results

Our main theoretical result is

Theorem 1 For $\delta \downarrow 0$ and $r \geq 1/2$, under conditions (V) and (M),

$$\mu_r^{-1} \delta^{1-r/2} [y_\delta^*]^{[r]}(t) \xrightarrow{p} \tau^{r/2*}(t), \quad (5)$$

and

$$\frac{\mu_r^{-1} \delta^{1-r/2} [y_\delta^*]^{[r]}(t) - \tau^{r/2*}(t)}{\mu_r^{-1} \delta^{1-r/2} \sqrt{\mu_{2r}^{-1} v_r [y_\delta^*]^{[2r]}(t)}} \xrightarrow{\mathcal{L}} N(0, 1), \quad (6)$$

²The equivalence follows on noting that for each j there exists an ω_j with

$$\inf_{(j-1)\delta \leq s \leq j\delta} \tau(s) \leq \omega_j \leq \sup_{(j-1)\delta \leq s \leq j\delta} \tau(s)$$

such that

$$|\tau^r(\eta_j) - \tau^r(\xi_j)| = r\omega_j^{r-1} |\tau(\eta_j) - \tau(\xi_j)|$$

and then using that τ is pathwise bounded away from 0 and ∞ .

³This condition is implied by Lipschitz continuity and itself implies continuity of α .

⁴Condition (V) is satisfied in particular if τ is of OU type and condition (M) is valid if, for instance, α is a intOU process plus drift.

where $\mu_r = E\{|u|^r\}$ and $v_r = \text{Var}\{|u|^r\}$, with $u \sim N(0, 1)$. \square

This theorem tells us that, for $\delta \downarrow 0$, scaled realised power variation converges in probability to integrated power volatility and follows asymptotically a normal variance mixture distribution with variance distributed as

$$\delta \mu_r^{-2} v_r \tau^{r*}(t),$$

which is consistently estimated by the square of the denominator in (6). Hence the limit theory is statistically feasible and does not depend upon knowledge of α^* or σ^2 .

Leading cases are *realised quadratic variation*, which is usually called *realised volatility* in the finance and econometrics literature,

$$[y_\delta^*]^{[2]}(t) = \sum_{j=1}^M y_j^2(t),$$

in which case

$$\frac{\sum_{j=1}^M y_j^2(t) - \tau^*(t)}{\sqrt{\frac{2}{3} \sum_{j=1}^M y_j^4(t)}} \xrightarrow{\mathcal{L}} N(0, 1), \quad (7)$$

and *realised absolute variation*

$$[y_\delta^*]^{[1]}(t) = \sum_{j=1}^M |y_j(t)|,$$

when

$$\frac{\sqrt{\pi/2} \sqrt{\delta} \sum_{j=1}^M |y_j(t)| - \sigma^*(t)}{\sqrt{(\pi/2 - 1) \delta \sum_{j=1}^M y_j^2(t)}} \xrightarrow{\mathcal{L}} N(0, 1). \quad (8)$$

In the case of $r = 2$ the result considerably strengthens the well known quadratic variation result that realised quadratic variation converges in probability to integrated volatility $\int_0^t \sigma^2(s) ds$ — which was highlighted in concurrent and independent work by Andersen and Bollerslev (1998a) and Barndorff-Nielsen and Shephard (2001). The asymptotic distribution of realised quadratic variation was discussed by Barndorff-Nielsen and Shephard (2002) in the special case where $\alpha^*(t) = \mu t + \beta \int_0^t \sigma^2(s) ds$. To our knowledge the probability limit of (normalised) realised absolute variation has not been previously derived, let alone its asymptotic distribution.

If on the left hand side of the main result (6) we standardised by $\delta^{(1-r/2)}$ rather than by $\mu_r^{-1} \delta^{1-r/2} \sqrt{\mu_{2r}^{-1} v_r [y_\delta^*]^{[2r]}(t)}$ we would obtain a normal variance mixture as the limit law, as follows from formula (5). Similar normal mixture results are known for standardised functions of the increments of diffusion processes, see Delattre and Jacod (1997) and Florens-Zmirou (1993). We also wish to point out that inspection of the proof of Proposition 1 in Section 4 below shows that the joint distribution of

$$\frac{\mu_r^{-1} \delta^{1-r/2} [y_\delta^*]^{[r]}(t) - \tau^{r/2*}(t)}{\mu_r^{-1} \delta^{1-r/2} \sqrt{\mu_{2r}^{-1} v_r [y_\delta^*]^{[2r]}(t)}}, \mu_{2r}^{-1} \delta^{1-r} [y_\delta^*]^{[2r]}(t)$$

converges to the product of $N(0, 1)$ and the law of an independent random variable distributed as $\tau^*(t)$.

Taking sums of squares of increments of log-prices has a very long tradition in financial economics — see, for example, Poterba and Summers (1986), Schwert (1989), Taylor and Xu (1997), Christensen and Prabhala (1998), Dacorogna, Muller, Olsen, and Pictet (1998), Andersen, Bollerslev, Diebold, and Labys (2001) and Andersen, Bollerslev, Diebold, and Ebens (2001). However, for a long time no theory was known for the behaviour of such sums outside the Brownian motion case. Since the link to quadratic variation has been made there has been a remarkably fast development in this field. Contributions include Corsi, Zumbach, Muller, and Dacorogna (2001), Andersen, Bollerslev, Diebold, and Labys (2001), Andersen, Bollerslev, Diebold, and Ebens (2001), Barndorff-Nielsen and Shephard (2002), Andreou and Ghysels (2001), Bai, Russell, and Tiao (2000), Maheu and McCurdy (2002), Areal and Taylor (2002), Galbraith and Zinde-Walsh (2000), Bollerslev and Zhou (2002) and Bollerslev and Forsberg (2002).

Andersen and Bollerslev (1998b) and Andersen and Bollerslev (1997) empirically studied the properties of $\sum_{j=1}^M |y_j(t)|$ computed using sums of absolute values of intra-day returns on speculative assets (many authors in finance have based their empirical analysis on absolute values of returns — see, for example, Taylor (1986, Ch. 2), Cao and Tsay (1992), Ding, Granger, and Engle (1993), West and Cho (1995), Granger and Ding (1995), Jorion (1995), Shiryaev (1999, Ch. IV) and Granger and Sin (2002)). This was empirically attractive, for using absolute values is less sensitive to possible large movements in high frequency data. There is evidence that if returns do not possess fourth moments then using absolute values rather than squares would be more reliable (see, for example, the work on the distributional behaviour of the correlogram of squared returns by Davis and Mikosch (1998) and Mikosch and Starica (2000)). However, the approach was abandoned in their subsequent work reported in Andersen and Bollerslev (1998a), Andersen, Bollerslev, Diebold, and Ebens (2001) and Andersen, Bollerslev, Diebold, and Labys (2001) due to the lack of appropriate theory for the sum of absolute returns as $\delta \downarrow 0$, although recently Andreou and Ghysels (2001) have performed some interesting Monte Carlo studies in this context, while Shiryaev (1999, pp. 349–350) and Maheswaran and Sims (1993) mention interests in the limit of sums of absolute returns. Our work provides, in particular, a theory for the use of sums of absolute returns.

4 Proofs

Since the mean and volatility processes α^* and τ are jointly independent of the Brownian motion w we need only argue conditionally on (α^*, τ) . Define α_j , τ_j and σ_j by

$$\alpha_j = \alpha^*(j\delta) - \alpha^*((j-1)\delta),$$

$$\tau_j = \tau^*(j\delta) - \tau^*((j-1)\delta)$$

and

$$\sigma_j = \sqrt{\tau_j}.$$

As a preliminary step we show

Lemma 1 For $\delta \rightarrow 0$,

$$\delta^{1-r} [\tau_\delta^*]^{[r]}(t) \rightarrow \tau^{r*}(t), \quad (9)$$

pathwise. \square

PROOF By the boundedness of $\tau(t)$ we have that for every $j = 1, \dots, M$ there exists a constant θ_j such that

$$\inf_{(j-1)\delta \leq s \leq j\delta} \tau(s) \leq \theta_j \leq \sup_{(j-1)\delta \leq s \leq j\delta} \tau(s)$$

and

$$\tau_j = \theta_j \delta, \quad (10)$$

and using this and the Riemann integrability of $\tau^r(t)$ we obtain

$$\begin{aligned} \delta^{1-r} [\tau_\delta^*]^{[r]} &= \delta^{1-r} \sum_{j=1}^M \tau_j^r \\ &= \sum_{j=1}^M (\tau_j/\delta)^r \delta \\ &= \sum_{j=1}^M \theta_j^r \delta \\ &\rightarrow \int_0^t \tau^r(s) ds \\ &= \tau^{r*}(t). \end{aligned}$$

\square

It is now convenient to write $y^*(t)$ as

$$y^*(t) = \alpha^*(t) + y_0^*(t),$$

where

$$y_0^*(t) = \int_0^t \sigma(s)dw(s)$$

and to introduce

$$y_{0j} = y_0^*(j\delta) - y_0^*((j-1)\delta).$$

The joint law of y_{01}, \dots, y_{0M} is equal to that of v_1, \dots, v_M where

$$v_j = \sigma_j u_j$$

and u_1, \dots, u_M are i.i.d. standard normal and independent of the process σ . It follows that

$$[y_{0\delta}^*]^{[r]}(t) \stackrel{\mathcal{L}}{=} \sum_{j=1}^M \tau_j^{r/2} |u_j|^r. \quad (11)$$

The conditional mean and variance of $[y_{0\delta}^*]^{[r]}(t)$ are then

$$\mathbb{E}\{[y_{0\delta}^*]^{[r]}(t)|\tau\} = \mu_r \sum_{j=1}^M \tau_j^{r/2} = \mu_r [\tau_\delta^*]^{[r/2]}(t) \quad (12)$$

and

$$\text{Var}\{[y_{0\delta}^*]^{[r]}(t)|\tau\} = v_r \sum_{j=1}^M \tau_j^r = v_r [\tau_\delta^*]^{[r]}(t). \quad (13)$$

Hence, letting

$$D_0 = \mu_r^{-1} [y_{0\delta}^*]^{[r]}(t) - [\tau_\delta^*]^{[r/2]}(t)$$

we have

$$\mathbb{E}\{D_0|\tau\} = 0$$

and

$$\text{Var}\{D_0|\tau\} = \mu_r^{-2} v_r [\tau_\delta^*]^{[r]}(t).$$

By Lemma 1 as $\delta \rightarrow 0$,

$$\delta^{1-r} \text{Var}\{D_0|\tau\} \rightarrow \mu_r^{-2} v_r \tau^{r^*}(t)$$

indicating the validity of

Proposition 1 For $\delta \rightarrow 0$,

$$\delta^{(1-r)/2} \frac{\mu_r^{-1} [y_{0\delta}^*]^{[r]}(t) - [\tau_\delta^*]^{[r/2]}(t)}{\sqrt{\mu_r^{-2} v_r \tau^{r^*}(t)}} \xrightarrow{\mathcal{L}} N(0, 1). \quad (14)$$

□

PROOF To establish this proposition we recall Taylor's formula with remainder term:

$$f(x) = f(0) + f'(0)x + x^2 \int_0^1 (1-s)f''(sx)ds. \quad (15)$$

By (11) and (12) we find, using (15), that the conditional cumulant transform of D_0 is of the form

$$\begin{aligned} \log E\{\exp(i\zeta D_0)|\tau\} &= \zeta^2 \sum_{j=1}^M \tau_j^r \int_0^1 (1-s)\kappa_r''(\tau_j^{r/2}\zeta s)ds \\ &= \zeta^2 \delta^r \sum_{j=1}^M \theta_j^r \int_0^1 (1-s)\kappa_r''(\delta^{r/2}\theta_j^{r/2}\zeta s)ds, \end{aligned}$$

where θ_j is given by (10) and κ_r denotes the cumulant transform of $\mu_r^{-1}|u|^r$ for u a standard normal random variable. Consequently,

$$\log E\{\exp(i\zeta\delta^{(1-r)/2}D_0)|\tau\} = \frac{1}{2}\zeta^2\delta R$$

where

$$R = 2 \sum_{j=1}^M \theta_j^r \int_0^1 (1-s)\kappa_r''(\delta^{1/2}\theta_j^{r/2}\zeta s)ds.$$

The boundedness of τ on $[0, t]$ implies

$$\lim_{\delta \downarrow 0} \delta^{1/2} \max_j \theta_j^{r/2} = 0$$

and hence, for $\delta \downarrow 0$,

$$\begin{aligned} \delta R &\rightarrow 2 \int_0^1 (1-s)ds \kappa_r''(0) \lim_{\delta \downarrow 0} \sum_{j=1}^M \theta_j^r \delta \\ &= \kappa_r''(0)\tau^{r*}(t) = -\mu_r^{-2}v_r\tau^{r*}(t). \end{aligned}$$

Therefore

$$\log E\{\exp(i\zeta\delta^{(1-r)/2}D_0)|\tau\} = -\frac{1}{2}\zeta^2\mu_r^{-2}v_r\tau^{r*}(t) + o(1) \quad (16)$$

and Proposition 1 follows. □

Lemma 1 uses only the local boundedness and Riemann integrability of τ . Invoking condition **(V)** we may strengthen the result (9) as follows.

Lemma 2 Under condition **(V)** we have

$$\delta^{1-r}[\tau_\delta^*]^{[r]}(t) - \tau^{r*}(t) = o_p(\delta^{1/2}).$$

□

PROOF For each j there exists a number ψ_j such that

$$\inf_{(j-1)\delta \leq s \leq j\delta} \tau(s) \leq \psi_j \leq \sup_{(j-1)\delta \leq s \leq j\delta} \tau(s)$$

and

$$\int_{(j-1)\delta}^{j\delta} \tau^r(s) ds = \psi_j^r \delta. \quad (17)$$

Using this and (10) we find

$$\begin{aligned} \delta^{1-r} [\tau_\delta^*]^{[r]}(t) - \tau^{r*}(t) &= \delta^{1-r} \sum_{j=1}^M \tau_j^r - \int_0^t \tau^r(s) ds \\ &= \delta \sum_{j=1}^M (\theta_j^r - \psi_j^r), \end{aligned} \quad (18)$$

and the conclusion now follows from assumption **(V)**. □

Proposition 2 Under condition **(V)**, for $\delta \rightarrow 0$,

$$\frac{\delta^{1-r/2} \mu_r^{-1} [y_{0\delta}^*]^{[r]}(t) - \tau^{r/2*}(t)}{\delta^{1/2} \sqrt{\mu_r^{-2} v_r \tau^{r*}(t)}} \xrightarrow{\mathcal{L}} N(0, 1). \quad (19)$$

□

PROOF We may rewrite the left hand side of (14) as

$$\begin{aligned} \frac{\delta^{1-r/2} \mu_r^{-1} [y_{0\delta}^*]^{[r]}(t) - \delta^{1-r/2} [\tau_\delta^*]^{[r/2]}(t)}{\delta^{1/2} \sqrt{\mu_r^{-2} v_r \tau^{r*}(t)}} &= \frac{\delta^{1-r/2} \mu_r^{-1} [y_{0\delta}^*]^{[r]}(t) - \tau^{r/2*}(t)}{\delta^{1/2} \sqrt{\mu_r^{-2} v_r \tau^{r*}(t)}} \\ &\quad + \frac{\delta^{1-r/2} [\tau_\delta^*]^{[r/2]}(t) - \tau^{r/2*}(t)}{\delta^{1/2} \sqrt{\mu_r^{-2} v_r \tau^{r*}(t)}} \end{aligned}$$

and Lemma 2 then implies the result. □

As an immediate consequence of Proposition 2 we have

Corollary 1 Under condition **(V)**, for $\delta \rightarrow 0$,

$$\delta^{1-r} \mu_{2r}^{-1} [y_{0\delta}^*]^{[2r]}(t) \xrightarrow{P} \tau^{r*}(t).$$

□

In other words, when normalised, $[y_{0\delta}^*]^{[2r]}(t)$ provides a consistent estimate of $\tau^{r^*}(t)$. Combining this with (19) yields the conclusion of Theorem 1 for the special case where the mean process α^* is identically 0.

The remaining task is to show that, to the order concerned, α^* does not affect the asymptotic limit behaviour, provided conditions **(V)** and **(M)** hold. For this it suffices to show that, under **(V)** and **(M)**,

$$\delta^{(1-r)/2} \left\{ [y_\delta^*]^{[r]}(t) - [y_{0\delta}^*]^{[r]}(t) \right\} = o_p(1),$$

cf. Proposition 1.

We shall in fact prove the following stronger result.

Proposition 3 Under conditions **(V)** and **(M)**,

$$\delta^{-r/2} \left\{ [y_\delta^*]^{[r]}(t) - [y_{0\delta}^*]^{[r]}(t) \right\} = O_p(1).$$

□

PROOF Let

$$\underline{\tau} = \inf_{0 \leq s \leq t} \tau(s) \quad \text{and} \quad \bar{\tau} = \sup_{0 \leq s \leq t} \tau(s)$$

and

$$\gamma_j = \delta^{-1} \alpha_j$$

and note that (pathwise for (α^*, τ)), by assumption,

$$0 < \underline{\tau} \leq \bar{\tau} < \infty,$$

implying

$$0 < \min_j \theta_j \leq \max_j \theta_j < \infty,$$

while, due to assumption **(M)**, there exists a constant c for which

$$\max_j |\gamma_j| \leq c,$$

whatever the value of M .

We have

$$\begin{aligned} [y_\delta^*]^{[r]}(t) - [y_{0\delta}^*]^{[r]}(t) &= \sum_{j=1}^M (|\alpha_j + y_{0j}|^r - |y_{0j}|^r) \\ &= \sum_{j=1}^M \left(|\delta\gamma_j + \delta^{1/2}\theta_j^{1/2}u_j|^r - |\delta^{1/2}\theta_j^{1/2}u_j|^r \right) \\ &= \delta^{r/2} \sum_{j=1}^M \theta_j^{r/2} \left\{ \left| \left(\gamma_j / \theta_j^{1/2} \right) \delta^{1/2} + u_j \right|^r - |u_j|^r \right\} \end{aligned}$$

and hence

$$\delta^{-r/2} \left\{ [y_\delta^*]^{[r]}(t) - [y_{0\delta}^*]^{[r]}(t) \right\} \stackrel{\mathcal{L}}{=} \sum_{j=1}^M \theta_j^{r/2} h_r \left(u_{0j}; \gamma_j / \theta_j^{1/2} \right)$$

where

$$h_r(u; \rho) = \left| \rho \delta^{1/2} + u \right|^r - |u|^r.$$

The conclusion of Proposition 3 now follows from Lemma 3 below. \square

Lemma 3 For $r \geq 1/2$, u a standard normal random variable and ρ constant,

$$\mathbb{E}\{h_r(u; \rho)\} = O(\delta)$$

and

$$\text{Var}\{h_r(u; \rho)\} = O(\delta).$$

\square

PROOF With φ denoting the standard normal density we obtain

$$\begin{aligned} \mathbb{E}\{|\rho \delta^{1/2} + u|^r\} &= \int_{-\infty}^{\infty} \left| \rho \delta^{1/2} + x \right|^r \varphi(x) dx \\ &= \int_{-\infty}^{\infty} |x|^r \varphi(x) e^{\rho \delta^{1/2} x} dx e^{-\rho^2 \delta / 2} \\ &= \int_{-\infty}^{\infty} |x|^r \varphi(x) e^{\rho \delta^{1/2} x} dx + O(\delta) \\ &= \mathbb{E}\{|u|^r\} + \int_{-\infty}^{\infty} |x|^r \varphi(x) \left(e^{\rho \delta^{1/2} x} - 1 \right) dx + O(\delta), \end{aligned}$$

i.e.

$$\begin{aligned} \mathbb{E}\{h_r(u; \rho)\} &= \delta^{1/2} \int_{-\infty}^{\infty} x |x|^r \varphi(x) \frac{e^{\rho \delta^{1/2} x} - 1}{\delta^{1/2} x} dx + O(\delta) \\ &= \delta^{1/2} \int_{-\infty}^{\infty} x |x|^r \varphi(x) \frac{e^{\rho \delta^{1/2} x} - 1 - \rho \delta^{1/2} x}{\delta^{1/2} x} dx + \rho \delta^{1/2} \int_{-\infty}^{\infty} x |x|^r \varphi(x) dx + O(\delta) \\ &= O(\delta). \end{aligned}$$

Furthermore,

$$\begin{aligned} \mathbb{E}\{h_r(u; \rho)^2\} &= \mathbb{E}\{h_{2r}(u; \rho)\} + 2\mathbb{E}\left\{|u|^r \left(|u|^r - |\rho \delta^{1/2} + u|^r \right)\right\} \\ &= O(\delta) + 2\mathbb{E}\left\{|u|^r \left(|u|^r - |\rho \delta^{1/2} + u|^r \right)\right\} \end{aligned} \tag{20}$$

by the previous result. Here

$$\mathbb{E}\{|u|^r |\rho \delta^{1/2} + u|^r\} = \int_{-\infty}^{\infty} |x|^r |\rho \delta^{1/2} + x|^r \varphi(x) dx$$

$$\begin{aligned}
&= \int_{-\infty}^{\infty} \left| x^2 - \frac{1}{4}\rho^2\delta \right|^r \varphi(x) e^{\frac{1}{2}\rho\delta^{1/2}x} dx e^{-\frac{1}{8}\rho^2\delta} \\
&= \int_{-\infty}^{\infty} \left| x^2 - \frac{1}{4}\rho^2\delta \right|^r \varphi(x) e^{\frac{1}{2}\rho\delta^{1/2}x} dx + O(\delta) \\
&= \int_{-\infty}^{\infty} |x|^{2r} \varphi(x) e^{\frac{1}{2}\rho\delta^{1/2}x} dx + O(\delta) \\
&\quad + \int_{-\infty}^{\infty} \left(\left| x^2 - \frac{1}{4}\rho^2\delta \right|^r - |x^2|^r \right) \varphi(x) e^{\frac{1}{2}\rho\delta^{1/2}x} dx. \tag{21}
\end{aligned}$$

Now, for a and b nonnegative numbers we have the inequality

$$|b^r - |b - a|^r| \leq \begin{cases} a^r & \text{for } 0 \leq b \leq a \\ rb^{r-1}a & \text{for } b > a. \end{cases} \tag{22}$$

Using this and $r \geq 1/2$ we find that

$$\begin{aligned}
&\left| \int_{-\infty}^{\infty} \left(\left| x^2 - \frac{1}{4}\rho^2\delta \right|^r - |x^2|^r \right) \varphi(x) e^{\frac{1}{2}\rho\delta^{1/2}x} dx \right| \\
&\leq \left(\frac{r}{4}\rho^2\delta \right)^r \int_{|x| \leq \rho\delta^{1/2}/2} \varphi(x) e^{\frac{1}{2}\rho\delta^{1/2}x} dx + \frac{r}{4}\rho^2\delta \int_{-\infty}^{\infty} |x|^{2(r-1)} \varphi(x) e^{\frac{1}{2}\rho\delta^{1/2}x} dx \tag{23} \\
&= O(\delta). \tag{24}
\end{aligned}$$

Thus, combining (20), (21) and (24), we have

$$\begin{aligned}
\mathbb{E}\{h_r(u; \rho)^2\} &= 2 \left\{ \int_{-\infty}^{\infty} |x|^{2r} \varphi(x) dx - \int_{-\infty}^{\infty} |x|^{2r} \varphi(x) e^{\frac{1}{2}\rho\delta^{1/2}x} dx \right\} + O(\delta) \\
&= 2 \int_{-\infty}^{\infty} |x|^{2r} \varphi(x) \left(1 - e^{\frac{1}{2}\rho\delta^{1/2}x} \right) dx + O(\delta) \\
&= 2\delta^{1/2} \int_{-\infty}^{\infty} x |x|^{2r} \varphi(x) \frac{1 - e^{\frac{1}{2}\rho\delta^{1/2}x}}{\delta^{1/2}x} dx + O(\delta) \\
&= O(\delta),
\end{aligned}$$

as was to be shown. \square

5 Examples

The following two examples show that conditions **(V)** and **(M)** are satisfied for OU models used by Barndorff-Nielsen and Shephard (2001) in the context of SV models.

Example 1 *Volatility process τ of OU type.* Without loss of generality we suppose that τ satisfies the equation

$$\tau(t) = e^{-\lambda t} \tau(0) + \int_0^t e^{-\lambda(t-s)} dz(\lambda s),$$

where z is a subordinator, i.e. a positive Lévy process, which is referred to as the BDLP (background driving Lévy process).

The number of jumps of the BDPL z on the interval $[0, t]$ is at most countable. Let $z_1 \geq z_2 \geq \dots$ denote the jump sizes given in decreasing order and let u_1, u_2, \dots be the corresponding jump times. Then, for any $s \in [0, t]$,

$$\tau(s) = \tau(0)e^{-\lambda s} + \sum_{n=1}^{\infty} z_n e(s; u_n),$$

where

$$e(s; u) = e^{-\lambda(s-u)} \mathbf{1}_{[u, t]}(s)$$

and we have

$$\sum_{j=0}^M |e(j\delta; u) - e((j-1)\delta; u)| \leq 2.$$

Hence

$$\sum_{j=0}^M |\tau(j\delta) - \tau((j-1)\delta)| \leq 2 \left\{ \tau(0) + \sum_{n=1}^{\infty} z_n \right\} = 2 \{ \tau(0) + z(\lambda t) \},$$

showing that condition **(V)** is amply satisfied. \square

Example 2 *OU volatility and intOU risk premium* In this particular case the volatility process τ is as in the previous example and the mean process is of the form

$$\alpha^*(t) = \mu\delta + \beta\tau^*(t),$$

where μ and β are arbitrary real parameters.

We then have

$$\alpha_j = \delta\{\mu + \beta\theta_j\},$$

implying

$$\max_{1 \leq j \leq M} \delta^{-1} |\alpha^*(j\delta) - \alpha^*((j-1)\delta)| \leq |\mu| + |\beta|\bar{\tau} < \infty,$$

so that condition **(M)** is indeed satisfied. \square

6 A Monte Carlo experiment

6.1 Multiple realised power variations

We have stated the definition and results for realised power variation for a single fixed t . It is clear that the theory can also be applied repeatedly on non-overlapping increments to the process. Let us write $\Delta > 0$ as a time interval and focus on the n -th such interval. Then define the intra- Δ increments as

$$y_{j,n} = y^* \left((n-1)\Delta + \frac{\Delta j}{M} \right) - y^* \left((n-1)\Delta + \frac{\Delta(j-1)}{M} \right), \quad \delta = \Delta M^{-1},$$

which allows us to construct the n -th realised power variation

$$[y_\delta^*]_n^{[r]} = \sum_{j=1}^M |y_{j,n}|^r.$$

The implication is that

$$\frac{\left[\mu_r^{-1} (\Delta M^{-1})^{1-r/2} [y_\delta^*]_n^{[r]} - \int_{(n-1)\Delta}^{n\Delta} \sigma^r(s) ds \right]}{\mu_r^{-1} (\Delta M^{-1})^{1-r/2} \sqrt{\frac{v_r}{\mu_{2r}} [y_\delta^*]_n^{[2r]}}} \xrightarrow{\mathcal{L}} N(0, 1).$$

An important observation is that the *realised power variation errors*

$$\left\{ \mu_r^{-1} (\Delta M^{-1})^{1-r/2} [y_\delta^*]_n^{[r]} \right\} - \sigma_n^{[r]}, \quad (25)$$

where $\sigma_n^{[r]}$, the *actual power volatility*, is defined as

$$\sigma_n^{[r]} = \sigma^{r*}(n\Delta) - \sigma^{r*}((n-1)\Delta), \quad \text{where} \quad \sigma^{r*}(t) = \int_0^t \sigma^r(s) ds,$$

will be asymptotically uncorrelated through n , although they will not be independent.

6.2 Simulated example

6.2.1 Realised power variation and actual power volatility

The above distribution theory says in particular that realised power variation error will converge in probability to zero as $\delta \downarrow 0$. To see the magnitude of this error we have carried out a simulation. This will allow us to see how accurate our asymptotic analysis is in practice. Throughout we have set the mean process $\alpha^*(t)$ to zero. Our experiments could have been based on the familiar constant elasticity of variance (CEV) process which is the solution to the SDE

$$d\sigma^2(t) = -\lambda \{ \sigma^2(t) - \xi \} dt + \omega \sigma(t)^\eta db(\lambda t), \quad \eta \in [1, 2],$$

where b is standard Brownian motion uncorrelated with w . Of course the special cases of $\eta = 1$ delivers the square root process, while when $\eta = 2$ we have Nelson's GARCH diffusion. These models have been heavily favoured by Meddahi and Renault (2002) in the context of SV models. Instead of this we will work with the non-Gaussian Ornstein-Uhlenbeck process, or OU process for short, which is the solution to the stochastic differential equation

$$d\sigma^2(t) = -\lambda \sigma^2(t) dt + dz(\lambda t), \quad (26)$$

where z is a subordinator cf. Examples 1 and 2 above. These models have been developed in this context by Barndorff-Nielsen and Shephard (2001). In Figure 1(a), (c), (e) we have drawn a curve to represent a simulated sample path of $\sigma_n^{[2]}$ from an OU process where $z(t)$ has a $\Gamma(t4, 8)$




Figure 1: *Simulation from an $OU-\Gamma(4,8)$ process. Drawn is $\sigma_n^{[2]}$ and $\sigma_n^{[1]}$ against time, together with their associated realised power variation estimators. Graph is computed for $M = 12, 48$ and 144 .*

marginal distribution, $\lambda = -\log(0.99)$ and $\Delta = 1$, along with the associated realised quadratic variation (depicted using crosses) computed using a variety of values of M . It is helpful to keep in mind that

$$\mathbb{E}(\sigma^2(t)) = \mathbb{E}(z(1)) = \frac{1}{2} \quad \text{and} \quad \text{Var}(\sigma^2(t)) = \frac{1}{2} \text{Var}(z(1)) = \frac{1}{32}.$$

The corresponding results for $\sigma_n^{[1]}$ and realised absolute variation is given in Figure 1(b), (d), (f). We see that as M increases the precision of realised power variation increases, while Figure 1 shows that the variance of the realised power variation increases with the level of volatility. This is line with the prediction from the asymptotic theory, for the denominator increases with the level of volatility.

6.2.2 QQ plots

To assess the finite sample performance of the asymptotic distributions of the realised quadratic and absolute variation we have constructed some QQ plots based upon the standardised errors (7) and (8). Our main focus will be on the absolute variation case, leaving the quadratic case to be covered in detail in a follow up paper by Barndorff-Nielsen and Shephard (2003). However, to start off with we give both cases, in order to allow an easy comparison.




Figure 2: Plots for the realised quadratic variation error and the realised absolute variation error plus twice their asymptotic standard errors. (a) and (b) have $M = 12$, while (c) and (d) has $M = 48$ and while (e) and (f) has $M = 144$. Corresponding QQ plots are on the right hand side, based on the standardised realised quadratic variations and the realised absolute variations.

Figure 2 gives QQ plots based on the simulation experiment reported in the previous subsection, with a sample size of 10,000. Again we vary M over 12, 48 and 144. The left hand side graphs give the realised quadratic variation errors $[y_\delta^*]_n^{[2]} - \sigma_n^{[2]}$ as well as plus and minus two times the asymptotic standard errors. The plot is based on the first 600 simulations. This graph

shows how much the standard errors change through time. This continues to happen with large values of M and reflects the stochastic denominator in the limit theory.

The middle graphs of Figure 2 give the corresponding realised absolute variation errors

$$\sqrt{\frac{\pi\delta}{2}} \sum_{j=1}^M |y_{j,n}| - \sigma_n^{[1]},$$

together with twice standard error bounds. The conditional standard errors are more stable through time, especially when M is small. The unconditional variance of the errors is approximately

$$\delta (\mu_1^{-2} - 1) \Delta E \{ \sigma^2(t) \}.$$

The right hand side of Figure 2 gives the associated QQ plots for the standardised residuals from the realised quadratic and absolute variation measures. These use all 10,000 observations. The results are clear, in comparison with the asymptotic limit laws both random variables are too fat tailed in small samples, with this problem reducing as M increases. The realised absolute variation version of the statistic has much better finite sample behaviour, while the realised quadratic variation is quite poorly behaved.

6.2.3 Logarithmic transformation

The realised power variation $[y_\delta^*]^{[r]}(t)$ is the sum of non-negative items and so is non-negative. It would seem sensible to transform this variable to the real line in order to improve its finite sample performance. Hence we use the standard logarithmic transformation (that is for a consistent estimator $\hat{\theta}$ of θ we approximate $\log(\hat{\theta})$ by $\log(\theta) + (\hat{\theta} - \theta)/\theta$, hence the asymptotic distribution of $\hat{\theta} - \theta$ can be used to deduce the asymptotic distribution of $\log(\hat{\theta}) - \log(\theta)$). For the general realised power variation this implies

$$\frac{\log [\mu_r^{-1} \delta^{1-r/2} [y_\delta^*]^{[r]}(t)] - \log \int_0^t \sigma^r(s) ds}{\mu_r^{-1} \delta^{1-r/2} \sqrt{\frac{v_r}{\mu_{2r}} \frac{[y_\delta^*]^{[2r]}(t)}{[\mu_r^{-1} \delta^{1-r/2} [y_\delta^*]^{[r]}(t)]^2}}} \xrightarrow{\mathcal{L}} N(0, 1). \quad (27)$$

Referring to the denominator we should note that

$$\frac{\mu_{2r}^{-1} \delta^{1-r} [y_\delta^*]^{[2r]}(t)}{[\mu_r^{-1} \delta^{1-r/2} [y_\delta^*]^{[r]}(t)]^2} \xrightarrow{\mathcal{L}} \frac{\int_0^t \sigma^{2r}(s) ds}{\left\{ \int_0^t \sigma^r(s) ds \right\}^2} \geq 1,$$

by Jensen's inequality. In the realised absolute variation case (27) simplifies to

$$\frac{\log \left\{ \sqrt{\frac{\pi\delta}{2}} \sum_{j=1}^M |y_j(t)| \right\} - \log \int_0^t \sigma(s) ds}{\sqrt{\delta \left(\frac{\pi}{2} - 1 \right) \frac{\sum_{j=1}^M y_j^2(t)}{\left\{ \sqrt{\frac{\pi\delta}{2}} \sum_{j=1}^M |y_j(t)| \right\}^2}}} \xrightarrow{\mathcal{L}} N(0, 1). \quad (28)$$




Figure 3: *Plots for the log transform of the realised absolute variation. Left hand plots are the errors plus and minus twice their asymptotic standard errors. Corresponding QQ plots are on the right hand side, based on the standardised log realised absolute variations.*

Using the same simulation setup as that employed in the previous subsection, we plot in Figure 3 the log version of the realised absolute variation error

$$\log \left\{ \sqrt{\frac{\pi\delta}{2}} \sum_{j=1}^M |y_{j,n}| \right\} - \log \sigma_n^{[1]},$$

plus and minus twice the corresponding standard errors using (28). The standard errors have now stabilised, almost not moving with n . This is not surprising for t times

$$\frac{t^{-1} \int_0^t \sigma^2(s) ds}{\left\{ t^{-1} \int_0^t \sigma(s) ds \right\}^2}$$

is empirically very close to one for small values of t while for large t it converges to

$$\frac{\mathbb{E} \{ \sigma^2(t) \}}{[\mathbb{E} \{ \sigma(t) \}]^2},$$

so long as the volatility process is ergodic and the moments exist.

The corresponding QQ plots in Figure 3 have also improved, with the normality approximation being accurate even for moderate values of M . This result carries over to wider simulations we have conducted.

Barndorff-Nielsen and Shephard (2003) have studied the effect of taking log-transforms in the case of $r = 2$. That paper used simulation and some theory to show the beneficial effect of taking a log and comparing its effectiveness with other transformations.

7 Conclusions

This paper has introduced the idea of realised power variation, which generalises the concept of realised volatility. The asymptotic analysis we provide, for $\delta \downarrow 0$, represents a significant extension of the usual quadratic variation result. Further, we provide a limiting distribution theory which considerably strengthens the consistency result and allows us to understand the variability of the difference between the realised power variation and the actual power volatility.

We have seen that when we take a log transformation of the realised power variation then the finite sample performance of the asymptotic approximation to the distribution of this estimator improves and seems to be accurate even for moderate values of M .

Our motivation for the study reported in this paper came originally from mathematical finance and financial econometrics where volatility is a key object of study. However, stochastic models in the form of a ‘signal’ α^* plus a noise term e where e is (conditionally) Gaussian with a variance that varies from ‘site’ to ‘site’ are ubiquitous in the natural and technical sciences, and we believe that results similar to those discussed here will be of interest for applications in a variety of other fields, for instance in turbulence and in spatial statistics (in this connection see Guyon and Leon (1989)).

Finally, the thesis of Becker (1998), briefly referred to in the Introduction, consists in a comprehensive study of the limit behaviour as $M \rightarrow \infty$ of processes of the type

$$y_M(t) = \sum_{j=1}^{[Mt]} f \left[\frac{j-1}{M}, \frac{1}{\sqrt{M}} \left\{ x \left(\frac{j}{M} \right) - x \left(\frac{j-1}{M} \right) \right\} \right]$$

where f is function of two variables and x denotes a Brownian semimartingale (of a certain kind, see below). The diffusion case is especially important. Extensions to general continuous or purely discontinuous semimartingales and even combination of the two are presented. The thesis is partly based on an earlier report by Jacod (1992), see also Delattre and Jacod (1997) and Florens-Zmirou (1993). Both x and f may be multidimensional, and generalisations to

cases where not only the increment of x over the j -th interval but the whole trajectory over that interval occur in the second argument of f are also considered.

We shall not here attempt to indicate the precise results and the accompanying regularity conditions of Becker's thesis in any detail, but we wish to underline that the setting of his study is extremely general. Of immediate interest in connection with the present paper are his results when x is a Brownian semimartingale. More specifically, Becker considers the case where x is of the form

$$x(t) = \int_0^t c(s)ds + \int_0^t \sigma(s)dw(s)$$

where w is Brownian motion and c and σ are predictable and subject to restrictions on their variational behaviour. He shows, in particular, that $y_M(t)$ after a suitable centering converges to a stochastic process which is representable as a certain type of stochastic integral where the integration is with respect to a 'martingale-measure tangential to x '. A key point of our present work is that for the kind of functions f we consider, i.e. absolute powers, we are able to identify the limit behaviour as mixed Gaussian and, crucially for the statistical applicability, from this to establish a standard normal limit statement using random rescaling by observable scale factors.

8 Acknowledgements

This paper represents a replacement of the paper "Higher order variation and stochastic volatility models," which first appeared on 4th July 2001. It had the result that (25) converges to zero in the case of r being 2, 4, 6, The general asymptotic distribution result contained in (6) was first discussed in public on 11th August 2001 at the Market Microstructure Conference, Centre for Analytical Finance, Denmark. Comments by Tim Bollerslev, Svend Erik Graversen, Matthias Winkel and the referees have been very helpful.

Ole E. Barndorff-Nielsen's work is supported by CAF (www.caf.dk), which is funded by the Danish Social Science Research Council, and by MaPhySto (www.maphysto.dk), which is funded by the Danish National Research Foundation. Neil Shephard's research is supported by the UK's ESRC through the grant "Econometrics of trade-by-trade price dynamics," which is coded R00023839.

References

- Andersen, T. G. and T. Bollerslev (1997). Intraday periodicity and volatility persistence in financial markets. *Journal of Empirical Finance* 4, 115–158.
- Andersen, T. G. and T. Bollerslev (1998a). Answering the skeptics: yes, standard volatility

- models do provide accurate forecasts. *International Economic Review* 39, 885–905.
- Andersen, T. G. and T. Bollerslev (1998b). Deutsche mark-dollar volatility: intraday activity patterns, macroeconomic announcements, and longer run dependencies. *Journal of Finance* 53, 219–265.
- Andersen, T. G., T. Bollerslev, F. X. Diebold, and H. Ebens (2001). The distribution of realized stock return volatility. *Journal of Financial Economics* 61, 43–76.
- Andersen, T. G., T. Bollerslev, F. X. Diebold, and P. Labys (2001). The distribution of exchange rate volatility. *Journal of the American Statistical Association* 96, 42–55.
- Andreou, E. and E. Ghysels (2001). Rolling-sampling volatility estimators: some new theoretical, simulation and empirical results. *Journal of Business and Economic Statistics* 19. Forthcoming.
- Areal, N. M. P. C. and S. J. Taylor (2002). The realized volatility of FTSE-100 futures prices. *Journal of Futures Markets* 22. Forthcoming.
- Bai, X., J. R. Russell, and G. C. Tiao (2000). Beyond Merton’s utopia: effects of non-normality and dependence on the precision of variance estimates using high-frequency financial data. Unpublished paper: Graduate School of Business, University of Chicago.
- Barndorff-Nielsen, O. E. and N. Shephard (2001). Non-Gaussian Ornstein–Uhlenbeck-based models and some of their uses in financial economics (with discussion). *Journal of the Royal Statistical Society, Series B* 63, 167–241.
- Barndorff-Nielsen, O. E. and N. Shephard (2002). Econometric analysis of realised volatility and its use in estimating stochastic volatility models. *Journal of the Royal Statistical Society, Series B* 64, 253–280.
- Barndorff-Nielsen, O. E. and N. Shephard (2003). How accurate is the asymptotic approximation to the distribution of realised volatility? In D. Andrews, J. Powell, P. A. Ruud, and J. H. Stock (Eds.), *Identification and Inference for Econometric Models. A Festschrift in Honour of Thomas J. Rothenberg*, Econometric Society Monograph Series. Cambridge: Cambridge University Press. Forthcoming.
- Becker, E. (1998). Theorems limite pour des processus discretises. These de Doctorat de l’Universite Paris 6.
- Bollerslev, T. and L. Forsberg (2002). The distribution of realised volatility and the normal inverse gaussian GARCH model: An application to the ECU and EURO exchange rates. *Journal of Applied Econometrics*. Forthcoming.

- Bollerslev, T. and H. Zhou (2002). Estimating stochastic volatility diffusion using conditional moments of integrated volatility. *Journal of Econometrics* 109, 33–65.
- Cao, C. Q. and R. S. Tsay (1992). Nonlinear time-series analysis of stock volatilities. *Journal of Applied Econometrics* 7, S165–S185.
- Christensen, B. J. and N. R. Prabhala (1998). The relation between implied and realized volatility. *Journal of Financial Economics* 37, 125–150.
- Corsi, F., G. Zumbach, U. Muller, and M. Dacorogna (2001). Consistent high-precision volatility from high-frequency data. Unpublished paper: Olsen and Associates, Zurich.
- Dacorogna, M. M., U. A. Muller, R. B. Olsen, and O. V. Pictet (1998). Modelling short term volatility with GARCH and HARCH. In C. Dunis and B. Zhou (Eds.), *Nonlinear Modelling of High Frequency Financial Time Series*. Chichester: Wiley.
- Davis, R. A. and T. Mikosch (1998). The limit theory for the sample ACF of stationary process with heavy tails with applications to ARCH. *Annals of Statistics* 26, 2049–2080.
- Delattre, S. and J. Jacod (1997). A central limit theorem for normalized functions of the increments of a diffusion process in the presence of round off errors. *Bernoulli* 3, 1–28.
- Ding, Z., C. W. J. Granger, and R. F. Engle (1993). A long memory property of stock market returns and a new model. *Journal of Empirical Finance* 1, 83–106.
- Florens-Zmirou, D. (1993). On estimating the diffusion coefficient from discrete observations. *Journal of Applied Probability* 30, 790–804.
- Galbraith, J. W. and V. Zinde-Walsh (2000). Properties of estimates of daily GARCH parameters based on intra-day observations. Unpublished paper: Economics Department, McGill University.
- Ghysels, E., A. C. Harvey, and E. Renault (1996). Stochastic volatility. In C. R. Rao and G. S. Maddala (Eds.), *Statistical Methods in Finance*, pp. 119–191. Amsterdam: North-Holland.
- Granger, C. W. J. and Z. Ding (1995). Some properties of absolute returns, an alternative measure of risk. *Annals d’Economie et de Statistique* 40, 67–91.
- Granger, C. W. J. and C.-Y. Sin (2002). Modelling the absolute returns of different stock indices: exploring the forecastability of an alternative measure of risk. *Journal of Forecasting*. Forthcoming.
- Guyon, X. and J. Leon (1989). Convergence en loi des H-variation d’un processus Gaussien stationnaire sur \mathbb{R} . *Ann. Inst. Henri Poincaré* 25, 265–282.

- Heston, S. L. (1993). A closed-form solution for options with stochastic volatility, with applications to bond and currency options. *Review of Financial Studies* 6, 327–343.
- Hull, J. and A. White (1987). The pricing of options on assets with stochastic volatilities. *Journal of Finance* 42, 281–300.
- Jacod, J. (1992). Limit of random measures associated with the increments of a Brownian semimartingale. Publ. Lab. Probabilite N. 120, Paris 6.
- Jorion, P. (1995). Predicting volatility in the foreign exchange market. *Journal of Finance* 50, 507–528.
- Lyons, T. (1994). Differential equations driven by rough signals. I. An extension of an inequality by L.C.Young. *Mathematical Research Letters* 1, 451–464.
- Maheswaran, S. and C. A. Sims (1993). Empirical implications of arbitrage-free asset markets. In P. C. B. Phillips (Ed.), *Models, Methods and Applications of Econometrics*. Basil Blackwell.
- Maheu, J. M. and T. H. McCurdy (2002). Nonlinear features of realised FX volatility. *Review of Economics and Statistics* 83. Forthcoming.
- Meddahi, N. and E. Renault (2002). Temporal aggregation of volatility models. *Journal of Econometrics*. Forthcoming.
- Mikosch, T. and R. Norvaiša (2000). Stochastic integral equations without probability. *Bernoulli* 6, 401–434.
- Mikosch, T. and C. Starica (2000). Limit theory for the sample autocorrelations and extremes of a GARCH(1,1) process. *Annals of Statistics* 28, 1427–1451.
- Nicolato, E. and E. Venardos (2001). Option pricing in stochastic volatility models of the Ornstein-Uhlenbeck type. *Mathematical Finance*. Forthcoming.
- Poterba, J. and L. Summers (1986). The persistence of volatility and stock market fluctuations. *American Economic Review* 76, 1124–1141.
- Renault, E. (1997). Econometric models of option pricing errors. In D. M. Kreps and K. F. Wallis (Eds.), *Advances in Economics and Econometrics: Theory and Applications*, pp. 223–78. Cambridge: Cambridge University Press.
- Schwert, G. W. (1989). Why does stock market volatility change over time? *Journal of Finance* 44, 1115–1153.
- Shiryaev, A. N. (1999). *Essentials of Stochastic Finance: Facts, Models and Theory*. Singapore: World Scientific.

Taylor, S. J. (1986). *Modelling Financial Time Series*. Chichester: John Wiley.

Taylor, S. J. and X. Xu (1997). The incremental volatility information in one million foreign exchange quotations. *Journal of Empirical Finance* 4, 317–340.

West, K. D. and D. Cho (1995). The predictive ability of several models for exchange rate volatility. *Journal of Econometrics* 69, 367–391.